



Stanford University

Non-Penetrating Schlemm's Canaloplasty versus Trabeculectomy

Diamond Y Tam* MD, Jennifer Calafati** MD, Iqbal Ike K. Ahmed**, MD, FRCSC



University of Toronto

*Dept. of Ophthalmology at Stanford University

**Dept. of Ophthalmology and Vision Sciences at the University of Toronto

Background


A trabeculectomy is a fistulizing procedure that allows aqueous humor to pass from the anterior chamber into the subconjunctival space. It is the currently accepted gold standard glaucoma surgery due to its efficacy in lowering intraocular pressure (IOP).

Canaloplasty is a non-penetrating surgical technique used for the treatment of open angle glaucomas which may be an alternative to conventional fistulizing trabeculectomy.

The mechanism of IOP lowering in canaloplasty is thought to be via augmentation of physiologic Schlemm's canal outflow, as well as through the intrascleral lake and into the episcleral/scleral venous plexi and suprachoroidal space. This obviates the need for a subconjunctival bleb, its inherent short and long term risks, and in theory lowers the risk of hypotony and its associated potentially visually devastating complications.

No study to date has reported a direct comparison of canaloplasty to trabeculectomy. Primary outcomes for the current study include IOP, number of glaucoma medications and BCVA at 6 and 12 months. Postoperative complications were examined as a secondary outcome.

Figure 1



Post-operative photograph of canaloplasty patient. Note the absence of a bleb

Objective

To compare and evaluate outcomes and complications after non-penetrating canaloplasty versus with conventional trabeculectomy as surgical treatment for patients with open angle glaucoma.

Methods



This study was a retrospective comparative chart review of 101 eyes of 93 patients. Fifty eyes underwent canaloplasty and 51 underwent trabeculectomy. Data indicating glaucoma severity (visual field indices and cup-to-disc ratio), IOP, number of glaucoma medications used, and best corrected visual acuity (BCVA) was collected. Time points were defined as preoperatively and after 1, 3, 6 and 12 months postoperatively. Complications and adjunctive procedures were recorded for each group. For a Kaplan-Meier survival analysis, complete success at 12 months was defined as a postoperative IOP of >6 mm Hg and <18 mm Hg without an increase in the number of glaucoma medications used postoperatively with no bleb needlings or further incisional procedures. Surgical or qualified success at 12 months was defined as an IOP of >6 and <18 mm Hg regardless of the number of medications used with no bleb needlings or further surgery. The use of postoperative laser suture lysis, laser goniopuncture, or transient hypotony was not considered a failure.

Results

IOP CONTROL

	IOP Control (mmHg)					
	Preop IOP	6 mo IOP	p-value*	Preop IOP	1 yr IOP	p-value*
Canaloplasty	26.4 ± 6.5	13.4 ± 3.3	<0.001	26.4 ± 6.5	13.4 ± 2.7	<0.001
Trabeculectomy	26.8 ± 8.1	12.5 ± 3.3	<0.001	26.8 ± 8.1	12.3 ± 3.5	<0.001
p-value**	0.803	0.172		0.803	0.103	

*IOP control within groups p-value
**IOP control between groups p-value



CP=Canaloplasty, TT=Trabeculectomy
*Complete success between groups p=0.313 (Log-Rank)
*Qualified success between groups p=0.010 (Log-Rank)

BCVA

	BCVA (LogMAR)					
	Preop VA	6 mo VA	p-value*	Preop VA	1 yr VA	p-value*
Canaloplasty	0.35 ± 0.58	0.25 ± 0.33	0.340	0.35 ± 0.58	0.29 ± 0.37	0.447
Trabeculectomy	0.50 ± 0.63	0.60 ± 0.82	0.301	0.50 ± 0.63	0.63 ± 0.88	0.192
p-value**	0.217	0.013		0.803	0.103	0.022


*BCVA within groups p-value
**BCVA between groups p-value

GLAUCOMA MEDICATIONS REQUIRED

	#GTTS					
	Preop gtts	6 mo gtts	p-value*	Preop gtts	1 yr gtts	p-value*
Canaloplasty	3.6 ± 0.9	0.7 ± 1.3	<0.001	3.6 ± 0.9	0.6 ± 1.1	<0.001
Trabeculectomy	3.6 ± 1.1	0.4 ± 0.9	<0.001	3.6 ± 1.1	0.7 ± 1.3	<0.001
p-value**	0.971	0.222		0.971	0.839	

*Average #GTTS within groups p-value
** Average #GTTS between groups p-value

Figure 2. Brief Summary of Steps in Canaloplasty



Results

Postoperative Complications

Complication	N, (%)		p-value*
	Canaloplasty	Trabeculectomy	
Choroidal effusion	1 (2)	14 (28)	<0.001
Transient hypotony (IOP ≤ 6 mmHg for at least 2 visits + resolved)	1 (2)	13 (26)	<0.001
Hyphema/microhyphema	9 (18)	9 (18)	0.482
Shallow/flat anterior chamber	0	8 (16)	0.002
Inadvertent bleb formation	11 (22)	N/A	-
Loss of >2 lines Snellen VA	1 (2)	8 (16)	0.008
Bleb fibrosis	0	6 (12)	0.006
IOP spike (increase in IOP ≥ 10 mmHg from one visit to next)	5 (10)	2 (4)	0.118
Wound leak	0	5 (10)	0.012
Bleb encapsulation	0	4 (8)	0.022
Cataract	3 (6)	3 (6)	0.490
Iris incarceration	3 (6)	2 (4)	0.317
Persistent hypotony (IOP ≤ 6 mmHg for at least 2 visits + resolve d)	1 (2)	2 (4)	0.286
Localized Descemet's detachment	2 (4)	0	0.080
Iris bombé	1 (2)	0	0.161
Malignant glaucoma	0	1 (2)	0.161
Bleb dyesthesia	0	1 (2)	0.161
Blebitis	0	1 (2)	0.161
Corneal decompensation	0	1 (2)	0.161
Cystic bleb	0	1 (2)	0.161
Epiretinal membrane	0	1 (2)	0.161
Persistent anterior chamber inflammation	1 (2)	0	0.161
Chronic cystoid macular edema	1 (2)	1 (2)	0.494
Retinal detachment	0	1 (2)	0.161
Late bleb leak	0	0	-
Endophthalmitis	0	0	-

*Between groups p-value



FIGURE 3. Postoperative Gonio-photograph of the Trabeculo-Descemet window created during canaloplasty with two 10-0 prolene suture knots visible.



FIGURE 4. Anterior segment OCT image showing centripetal indentation of the Schlemm's canal as a result of the intracanalicular tensioned suture.

Conclusion

- Non-penetrating Schlemm's canaloplasty and MMC trabeculectomy were found to have equal efficacy with regards to IOP lowering and decrease in glaucoma medication usage at up to 1 year postoperatively.
- Eyes undergoing canaloplasty recovered BCVA faster postoperatively than eyes undergoing trabeculectomy.
- Eyes undergoing canaloplasty sustained fewer complications than those undergoing trabeculectomy.

References

1. Cairns J. Trabeculectomy. Preliminary report of a new method. Am J Ophthalmol 1968;66:673-9.
2. Fyodorov S, Ioffe DI, Ronika TI. Deep sclerectomy: technique and mechanism of a new antiglaucomatous procedure. Glaucoma 1984;6:281-3.
3. Zimmerman T, Koener KS, Ford VJ, et al. Trabeculectomy vs. nonpenetrating trabeculectomy: a retrospective study of two procedures in phakic patients with glaucoma. Ophthalmic Surg 1984;15:734-40.
4. Lewis R, von Wolff K, Tetz M, et al. Canaloplasty: circumferential viscodilation and tensioning of Schlemm's canal using a flexible microcatheter for the treatment of open-angle glaucoma in adults: two-year interim clinical study results. J Cataract Refract Surg 2009 May;35:814-24.